arXiv:1006.5147v1 [gr-qc] 26 Jun 2010

On black holes in multidimensional theory with Ricci-flat internal spaces

S.B. Fadeev, V.D. Ivashchuk and V.N. Melnikov

Center for Gravitation and Fundamental Metrology, VNIIMS, 46 Ozyornaya Str., Moscow 119361, Russia [1]

Abstract

A generalization of the Tangherlini solution for the case of n internal Ricci-flat spaces is obtained. It is shown that in the $(2+d)$-dimensional section a horizon exists only in the trivial case when the internal-space factors are constant. The p-adic analog of the solution is also considered.

1. In ref. [1] the Schwarzschild solution is generalized for the case of n internal Ricci-flat spaces. It was shown that a horizon in the four-dimensional section of the metric exists only when the internal space scale factors are constant. This proposition was also suggested in ref. [2], where a special case of the solution [1] with $n = 2$ internal spaces (one of which being one dimensional) was considered.

In this paper we consider exact static, spherically symmetric solutions of the Einstein equations in $(2+d+N_1+\ldots+N_n)$-dimensional gravity $(d \geq 2)$ with a chain of n Ricci-flat internal spaces. We show that as in the $d = 2$ case, considered in ref. [1], a horizon is absent in all nontrivial cases. Finally, we consider a formal analog of the solution for the case of p-adic numbers [3].

2. We consider the Einstein equations

$$R_{MN} = 0 \tag{1}$$

on the D-dimensional manifold

$$M = M_0 \times M_1 \times \ldots \times M_n, \tag{2}$$

---

[1] the current address

where

$$\dim\ M_i = N_i, \quad D = 2 + d + \sum_{j=1}^{n} N_j, \quad i > 0,$$

M_0 is $(2+d)$-dimensional space-time $(d \geq 2)$ and M_i are Ricci-flat manifolds with the metrics $g_{(i)}, i = 1, \ldots, n$. We seek solutions of (1) such that M_0 is static, spherically symmetric ($O(d+1)$-symmetric), while all the scale factors $\exp(\beta_i)$ of the internal spaces M_i depend on the radial coordinate u, i.e., the D-metric is

$$\begin{aligned} g = & - \exp[2\gamma(u)]dt \otimes dt + \exp[2\alpha(u)]du \otimes du \\ & + \exp[2\beta(u)]d\Omega_d^2 + \sum_{i=1}^{n} \exp[2\beta_i(u)]g_{(i)}, \end{aligned} \tag{3}$$

where $d\Omega_d^2 = g_{(0)}$ is the standard S^d metric.

If we denote $\gamma = \beta_{-1}, N_{-1} = 1$ and $\beta = \beta_0, N_0 = d$ and choose the harmonic radial coordinate u such that $\alpha = \sum_{k=-1}^{n} \beta_i N_i$ then the Einstein equations (1) can be written in the form

$$\begin{aligned} R_{11} &= \sum_{i=-1}^{n} (-\beta_i^{''} + \alpha^{'}\beta_i^{'} - (\beta_i^{'})^2)N_i = 0, \\ R_{00} &= \exp(2\beta_{-1} - 2\alpha)\beta_{-1}^{''} = 0, \\ R_{kl} &= g_{(0)kl}[d - 1 - \beta_0^{''}\exp(2\beta_0 - 2\alpha)] = 0. \\ R_{m_i n_i} &= -g_{(i)m_i n_i}\beta_i^{''}\exp(2\beta_i - 2\alpha),\ i = 1, \ldots, n. \end{aligned} \tag{4}$$

This set of equations is easily solved, so that the metric g (3) after an appropriate redefinition of the radial coordinate $(u \Rightarrow R = R(u))$ may be written in the following way,

$$\begin{aligned} g &= -c^2[1 - \varepsilon(L/R)^{d-1}]^a dt \otimes dt \\ &+ [1 - \varepsilon(L/R)^{d-1}]^{(a+b+d-2)/(1-d)} dR \otimes dR \\ &+ [1 - \varepsilon(L/R)^{d-1}]^{(a+b-1)/(1-d)} R^2 d\Omega_d^2 \\ &+ \sum_{i=1}^{n} c_i [1 - \varepsilon(L/R)^{d-1}]^{a_i} g_{(i)}, \\ \varepsilon &= \pm 1, \end{aligned} \tag{5}$$

where $L \geq 0$, $R > 0$ and $R > L$ for $\varepsilon = +1$; L, $c \neq 0$ and $c_i \neq 0$ are constants,

$$b = \sum_{i=1}^{n} a_i N_i, \tag{6}$$

and the constants $a, a_1, \ldots, a_n$ satisfy the relation

$$(a+b)^2 + (d-1)\left(a^2 + \sum_{i=1}^{n} a_i^2 N_i\right) = d. \tag{7}$$

In the case $d = 2$ and $\varepsilon = +1$ this solution coincides with that of ref. [1]. In the special case $n = 1$ and

$$a = \left(\frac{d(d+N_1-1)}{(d-1)(d+N_1)}\right)^{1/2}, \qquad a_1 = -\frac{a}{d+N_1-1},$$

this solution was considered earlier in ref. [4]. The case $d = 2$, $n = 1$ was considered in ref. [5], and its scalar-vacuum generalization was obtained in ref. [6].

3. Let us consider the $(2+d)$-dimensional section of the metric (5). In the case $L = 0$ the metric is flat, while for $L > 0$ and

$$a - 1 = a_1 = \ldots = a_n = 0 \tag{8}$$

it coincides with the Tangherlini solution [7].

Now let us prove that a horizon at $R = L$ ($L > 0$) takes place only in the case (8) for $\varepsilon = +1$. Indeed, for the light propagating along a radius from a place with $R = R_0$ towards the center the coordinate time interval is

$$t - t_0 = \frac{1}{c}\int_{R}^{R_0} dx[1-(L/R)^{d-1}]^{\lambda}, \tag{9}$$

where

$$\lambda = \frac{1}{2}\left(\frac{a+b+d-2}{1-d} - a\right). \tag{10}$$

Relation (7) is equivalent to the identity

$$(a+b/d)^2 = 1 - \frac{d-1}{d}\sum_{i=1}^{n} a_i^2 N_i - \frac{b^2}{d^2}(d-1). \tag{11}$$

Let $\varepsilon = +1$. If some a_i ($i = 1, \ldots, n$) are nonzero, then by (11) $a + b/d < 1$ and from (10) $\lambda > -1$, hence, the integral (9) converges at $R = L$. This means that a radial light beam reaches the surface $R = L$ in a finite time interval, i.e. it is not a horizon. When $a_i = 0$, $i = 1, \ldots, n$, then $a = \pm 1$. For the Tangherlini case $a = +1$ we have a horizon ($\lambda = -1$), and for $a = -1$ ($\lambda = 1/(d-1)$) the horizon at $R = L$ is absent. Evidently, for $\varepsilon = -1$ the horizon at $R = L$ is absent too. This completes the proof.

4. At present there is more interest in considering the physical models with p-adic numbers [3] instead of real ones. This interest was stimulated mainly by the pioneering works on p-adic strings [8, 9]. Recently a p-adic generalization of the classical and quantum gravitational theory was defined [10] and some solutions of the Einstein equations were considered [10, 11]. In this section we consider the p-adic analog of the solution (5). Let us briefly recall the definition of p-adic numbers [3, 12]. Let p be a prime number. Any rational number $a \neq 0$ can be represented in the form $a = p^k m/n$, where the integer numbers m and n are not divisible by p. Then the p-adic norm is defined as follows: $\mid a \mid_p = p^{-k}$. This norm is non-Archimedean: $\mid a + b \mid_p \leq \max(\mid a \mid_p, \mid b \mid_p)$. The completion of Q with this norm is the p-adic number field Q_p. Any nonzero p-adic number $a \in Q_p$ can be uniquely represented as the series

$$a = p^k (a_0 + a_1 p + a_2 p^2 + \ldots),$$

where $a_0 = 1, \ldots, p-1$, and $a_i = 0, \ldots, p-1$ for $i > 0$.

The definitions of derivatives, manifold and tensor analysis in the p-adic case are similar to those of the real case. The power p-adic function is defined as follows,

$$(1 + x)^{\alpha}_p \equiv \exp_p\{\alpha[\log_p(1 + x)]\}, \tag{12}$$

where $\mid x \mid_p < 1$ and $\mid \alpha \mid_p \mid x \mid_p < \delta_p$. Here $\delta_p = 1$ for $p \neq 2$ and $\delta_2 = \frac{1}{2}$. The definition is correct, for the functions $\exp_p$ and $\log_p$ are well defined on the discs $\{\mid x \mid_p < \delta_p\}$ and $\{\mid x - 1 \mid_p < 1\}$ respectively [3].

Let us consider the p-adic manifold

$$Q_p \times Q^*_p \times S^d \times M_1 \times \ldots \times M_n, \tag{13}$$

where $(S^d, g_{(0)})$ is a space of constant curvature

$$R^0_{ijkl} = g^{(0)}_{ik} g^{(0)}_{jl} - g^{(0)}_{il} g^{(0)}_{jk}$$

$[Q_p^* \subset Q_p]$ and $(M_i, g_{(i)})$ are Ricci-flat manifolds.

For $R \neq 0$ and

$$\left|\frac{L}{R}\right|_p^{d-1} < \min(1, 1/\mid a_i \mid_p), \quad i = -2, -1, \dots, n, \tag{14}$$

with

$$a_{-2} = \frac{a+b+d-2}{1-d}, \quad a_{-1} = a, \quad a_0 = \frac{a+b-1}{1-d},$$

the metric (5) on the manifold (13) is well defined. Then the Einstein equations for the metric (5), (14) on the manifold (13) are satisfied identically, when the parameters $a, a_1, \dots, a_n \in Q_p$ obey the restriction (7).

This can be easily checked using the identity

$$[(1+x)^{\alpha}]' = \alpha(1+x)^{\alpha}/(1+x),$$

$\mid x \mid_p, \mid \alpha \mid_p \mid x \mid_p < 1$ (the verification of (1) in the p-adic case is just the same as in the real one).

In the $d = 2$ case this solution was considered earlier in ref. [11]. It was pointed out that there is an infinite number or rational solutions of (7) in this case. For example, we may consider the set [11]

$$\begin{aligned} a_1 &= \frac{4k}{N_1(N_1+2)k^2+1}, \ a_i = 0, \ i > 1, \\ a &= \frac{-2N_1 k \pm [k^2 N_1(N_1+2) - 1]}{k^2 N_1(N_1+2)+1}, \ k \in \mathbb{Z}. \end{aligned}$$

In the p-adic case there exist pseudo-constant functions $C = C(R)$ such that $C'(R) = 0$ but $C(R)$ is not identically constant [12]. Such functions may be used in generalization of well-known solutions of differential equations. In our case there is also a possibility for the constants $c, c_1, \dots, c_n$ and $a, a_1, \dots, a_n$ to be replaced by the pseudo-constants (of course, the restriction (14) should be preserved).

There is another possibility to generalize the solution (5). We may suppose that the components of the metric g_{MN} belong to some extension of Q_p. It may be the quadratic extension of Q_p or even Ω_p, which is the completion of the algebraic closure of Q_p [3]. In this case the constants in (5) may belong to the extension of Q_p.

The solution (5) can be also generalized on scalar-vacuum and electrovacuum cases. The last generalization for $d = 2$ is considered in ref. [13].

References

[1] K.A. Bronnikov, V.D. Ivashchuk and V.N. Melnnikov, in: Problems of Gravitation, Plenary Reports of 7th. Soviet Conf. on Gravitation (ErGU, Erevan, 1989), p. 70.

[2] C.G. Callan, R.C. Myers and M.J. Perry., Nucl. Phys. B 311 (1988) 673.

[3] N. Koblitz, p-adic numbers, p-adic analysis and zeta-functions (Springer, Berlin, 1977).

[4] R.C. Myers, Phys. Rev. D 35 (1987) 455.

[5] M. Yoshimura, Phys. Rev. D 34 (1986) 1021.

[6] K.A. Bronnikov and V.D. Ivashchuk, in: Reports of 7th Soviet Conf. on Gravitation (ErGU, Erevan, 1988) p. 156-157 [in Russian].

[7] F.R. Tangherlini, Nuovo Cimento 27 (1963) 636.

[8] I.V. Volovich, Class. Quantum Grav. 4 (1987) 183.

[9] P.G.O. Freund and M. Olson, Phys. Lett. B 199 (1987) 186.

[10] I. Ya. Aref'eva, B. Dragovich, P. Frampton and I.V. Volovich, Wave function of the universe and p-adic gravity, Steklov Mathematical Institute Preprint (1990).

[11] V.D. Ivashchuk, V.N. Melnikov and S.B. Fadeev, in: Proc. National Workshop on Gravitation and Gauge Theory, Yakutsk (1990).

[12] K. Mahler, Introduction to p-adic numbers and their functions (Cambridge Univ. Press., Cambridge, 1973).

[13] S.B. Fadeev, V.D. Ivashchuk and V.N. Melnikov, Chinese Phys. Lett. (1991) [8 (1991) 439].